MVA PROCEDURE NOTE:

	cal Examination:
\sim .	s: Size in weeks (bimanual): AV/Mid / RV
	x: WNL / CMT, parous/nullip
Vagın	a: WNL / discharge noted:
D	A
	dure:
	Pap done gc, chlam done
	Cervix and vagina swabbed with Betadine.
	Lidocaine 1%,cc total injected. Tenaculum appliedo'clock.
	Tenaculum applied O clock.
	Cervix progressively dilated to:
□	Cannula inserted, size
	ated blood loss: cc.
Addit	ional comments:
Ti-	· Francisco
	e Exam:
	Decidual tissue
	Villi
	Gestational sac
Ц	Tissue appropriate for gestational age
D 44	
Post-G	Op Ultrasound if done:
Ц	No IUP visualized Other:
	sment:
	Patient stable, AB complete
Ш	Pad checked for bleeding
_	For complications, see progress notes
L	Post-procedure vital signs:
	B/P
<u>Plan:</u>	Expected symptoms discussed; post-procedure instructions given.
Plan: □	Rhogam if needed:
<u>Plan:</u> □	Rhogam if needed: Doxycycline 100 mg tabs 2 tab Dispensed or 1 gm Zithromax
<u>Plan:</u> □	Rhogam if needed: